arXiv:1612.03141v1 [math.AG] 9 Dec 2016

A NON-EXISTENCE RESULT FOR DE JONQUIÈRES DIVISORS

MARA UNGUREANU

Abstract. This paper represents a first step towards settling the issue of the validity of the de Jonquières formulas. We consider the space of divisors with prescribed multiplicity, or de Jonquières divisors, contained in a linear series on a smooth projective curve. Assuming zero expected dimension of this space, the de Jonquières formulas compute the virtual number of de Jonquières divisors. We show that if the expected dimension is negative, then a general linear series on a general curve does not admit de Jonquières divisors of the expected type.

1. Introduction

In his 1866 memoir [dJ], de Jonquières computed the number of divisors with prescribed multiplicities that are contained in a fixed linear series on a given plane algebraic curve.

Almost a century later, and using modern techniques of topology and intersection theory of cycles on the symmetric product on a curve, MacDonald [Mac] and Mattuck [Mat] recovered the original formula in characteristic zero and arbitrary characteristic, respectively and generalised it to linear series of any dimension. However, their work does not address the vagueness of the classical statements, assuming either that the linear series in question is sufficiently generic, or that the multiplicities are counted correctly. To address this issue, Vainsencher [Vai] described the locus of divisors with prescribed multiplicities as the vanishing locus of a section of a bundle of the appropriate rank. Using a natural filtration of this bundle, he computed its Chern classes without making use of the Grothendieck-Riemann-Roch theorem, and established the enumerative validity of the de Jonquières formula for plane curves and for some higher dimensional cases.

This paper represents a first step towards settling the issue of the validity of the de Jonquières formula for arbitrary r using a variational approach. In other words our aim is to specify the conditions under which linear series on a curve are allowed to contain a divisor with given multiplicities.

We fix the notation. Let C be a smooth projective curve of genus g and denote by C_d its d-th symmetric product and by $\mathrm{G}^{\mathrm{r}}_{\mathrm{d}}(\mathrm{C})$ its associated variety of linear series of type g^r_d, i.e.

$$\mathrm{G}^{\mathrm{r}}_{\mathrm{d}}(\mathrm{C}) := \{\mathcal{L} = (L, V) \mid L \in \mathrm{Pic}^d(C), V \in G(r+1, H^0(C, L))\}.$$

Brill-Noether theory shows that if $[C] \in \mathcal{M}_g$ is a general curve then $\mathrm{G}^{\mathrm{r}}_{\mathrm{d}}(\mathrm{C})$ is a smooth variety and its dimension is given by the Brill-Noether number

$$\rho(g, r, d) = g - (r+1)(g-d+r).$$

Such a linear series $\mathcal{L}$ on the curve C is said to admit a *de Jonquières divisor of length n* if

$$a_1p_1 + \ldots + a_np_n \in \mathbb{P}\,V$$

for some points $p_1, \ldots, p_n \in C$ and some positive integers $a_1, \ldots, a_n$ such that $a_1 + \ldots + a_n = d$. We call the corresponding divisor $D := p_1 + \ldots + p_n \in C_n$ a *de Jonquières divisor of length n* and we denote by $DJ_n(\mathcal{L})$ the space of de Jonquières divisors of length n corresponding to the linear series $\mathcal{L}$.

The de Jonquières formula (cf. [ACGH] chapter VIII, §5) states that, if we expect there to be a finite number of de Jonquières divisors of length n, then this virtual number is given by the coefficient of the monomial $t_1 \cdot \ldots \cdot t_n$ in

$$(1 + a_1^2t_1 + \ldots + a_n^2t_n)^g(1 + a_1t_1 + \ldots + a_nt_n)^{d-r-g}.$$

Our main result is the following

Theorem 1. *Let C be a general curve and consider a triple (n, r, d) of positive integers with $n - d + r < 0$. Then a general linear series $\mathcal{L} \in G^r_d(C)$ does not admit a de Jonquières divisor of length n. In other words, $DJ_n(\mathcal{L})$ is empty.*

We make a simple observation, which provides the first piece of evidence towards the validity of our result. Eisenbud and Harris prove in [EH] that a general linear series of any degree and dimension on a general curve may have only ordinary ramification. As a direct consequence, we can immediately exclude the existence of any de Jonquières divisors where $a_i \geq r+2$ for some $i \in \{1, 2, \ldots, n\}$. If this is the case, then clearly $n \leq d-r-1$, which agrees with our statement.

Note that Theorem 1 holds trivially for rational normal curves and normal elliptic curves. Indeed, in the case of the rational normal curve $n - d + r = n$, while for an elliptic curve $n - d + r = n - 1$. Clearly, if $n < 0$ or $n < 1$, respectively, there are no de Jonquières divisors of length n with positive degree, and hence no general linear series admitting de Jonquières divisors of length n.

In the particular case of the canonical linear series, our problem is naturally related to that of strata of abelian differentials which have recently received a lot of attention, for example in the work of Farkas and Pandharipande [FP] in the algebro-geometric context, or that of Bainbridge, Chen, Gendron, Grushevsky, and Möller [BCG$^+$] in the Teichmüller dynamics and flat surfaces context. In fact, the result of Polishchuk [Pol] concerning the dimension of the strata turns out to be a crucial ingredient in the proof of our theorem in the case of linear series with index of speciality $s = 1$.

The paper is organised as follows. In Section 2 we describe the space of de Jonquières divisors as a degeneracy locus of vector bundles and thus obtain a lower bound for its dimension, while Section 3 is dedicated to the proof of Theorem 1.

Acknowledgements

This paper is part of my PhD thesis. I would like to thank my advisor Gavril Farkas for introducing me to this circle of ideas and for his helpful suggestions to approaching this problem. I am also grateful to Daniele Agostini, Ignacio Barros, and especially Irfan Kadıköylü for many useful discussions.

2. The Space of de Jonquières Divisors

The space $DJ_n(\mathcal{L})$ can be described as a degeneracy locus of vector bundles on C_n as follows. The idea is that the condition

$$a_1p_1 + \ldots + a_np_n \in \mathbb{P}\,V$$

is equivalent to the condition that the natural restriction map

$$V \to H^0(C, L/L(-a_1p_1 - \ldots - a_np_n))$$

has non-zero kernel. These maps constitute a morphism $\phi : E \to F$ over C_n, where $E = \mathcal{O}_{C_n} \otimes V$ is the trivial bundle. To construct F, consider the diagram

$$\begin{array}{ccccc} & & C \times C_n \times C_d & \supset \Delta & \\ & \swarrow^{\sigma} & & \searrow^{\tau} & \\ C & & & & C_n \end{array}$$

where σ and τ are the usual projections and Δ is given by

$$\Delta := \{(p, p_1 + \ldots + p_n, D) \mid p \in C \text{ and } D = a_1p_1 + \ldots + a_np_n\} \subset C \times C_n \times C_d.$$

Set $F = \tau_*(\sigma^*L \otimes \Delta)$, which is indeed a vector bundle by the cohomology and base change theorem. Hence DJ_n is the locus in C_n where $\operatorname{rk}\phi \geq r$.

Lemma 2. *For every point $D \in DJ_n(\mathcal{L})$, one has $\dim_D DJ_n(\mathcal{L}) \geq n - d + r$.*

Proof. From the description of $DJ_n(\mathcal{L})$ as a degeneracy locus, we get that its codimension in C_n is at most

$$(\operatorname{rk} E - r)(\operatorname{rk} F - r) = (r + 1 - r)(d - r) = d - r. \qquad \square$$

3. The Case of Negative Expected Dimension

In this section we prove Theorem 1. The plan of the proof is as follows. The case $r = 1$ follows immediately from previously known results, which we summarise below. For $r \geq 2$, let $s := g - d + r$. We first prove the statement for non-special linear series (i.e. $s = 0$) via an easy argument using an Abel-Jacobi-type map and for linear series with $s = 1$ using a result of Polishchuk [Pol]. For the proof in the case of non-complete linear series (corresponding to $s < 0$) and in the case of series with $s > 1$ we use an induction argument based on degenerations of curves embedded in projective space.

3.1. **The case of general pencils.** A Kodaira-Spencer deformation theoretic argument (cf. [ACG] chapter XXI, §11) shows that the general point of any component of $\mathcal{W}^1_d$ corresponds to a smooth curve C together with a complete, base-point-free, simple g^1_d. This means that a general g^1_d on a general curve may only be of the form

$$\mathcal{O}(2p_1 + p_2 + \ldots + p_{d-1}),$$

which is equivalent to requiring that $n \geq d - 1$. Hence we conclude that if we have a triple $(n, 1, d)$ with $n - d + 1 < 0$, then the general linear series $\mathcal{L}$ of

degree d and dimension 1 on a general curve does not admit a de Jonquières divisor of length n.

3.2. **The case of non-special linear series of any dimension.**

Proposition 3. *Let C be an arbitrary curve of genus g and let (n, r, d) be a triple such that $n - d + r < 0$ and $s = 0$. Then a general linear series $\mathcal{L}$ of degree d and dimension r does not admit a de Jonquières divisor of length n. Moreover, for any $r' < r$, the general non-complete linear series of type $g^{r'}_d$ does not admit a de Jonquières divisor of length n either.*

Proof. Consider the following Abel-Jacobi-type map:

$$\begin{aligned} \varphi : C_n &\to \mathrm{Pic}^d(C) \\ (p_1, \ldots, p_n) &\mapsto \mathcal{O}(a_1 p_1 + \ldots + a_n p_n). \end{aligned}$$

Note that in the non-special regime $\mathrm{Pic}^d(C) = W^r_d(C)$. Moreover the image of φ is closed and $\dim \mathrm{im}\, \varphi \leq n < g = \dim \mathrm{Pic}^d(C)$. Thus a general line bundle $L \in \mathrm{Pic}^d(C)$ is not contained in the image of φ whence we conclude that the divisor $a_1 p_1 + \ldots + a_n p_n$ is not contained in a general linear series $\mathcal{L}$ of type g^r_d.

To prove the second part of the statement, let c be the forgetful morphism of varieties

$$\begin{aligned} c : G^{r'}_d &\to \mathrm{Pic}^d(C) \\ (L, V) &\mapsto L. \end{aligned}$$

Fix a complete non-special $L \in \mathrm{Pic}^d(C) \setminus \mathrm{im}\, \varphi$. It does not admit a de Jonquières divisor of length n, whence none of the linear series $l \in c^{-1}(L)$ will admit such a divisor either. Since c is continuous, $c^{-1}(\mathrm{Pic}^d(C) \setminus \mathrm{im}\, \varphi)$ is also open in $G^{r'}_d$ and clearly non-empty, and moreover no $l \in c^{-1}(\mathrm{Pic}^d(C) \setminus \mathrm{im}\, \varphi)$ admits a de Jonquières divisor of length n. Hence the general series of type $g^{r'}_d$ does not admit a de Jonquières divisor of length n. □

As seen in the second part of Proposition 3, the non-existence result for linear series of type g^r_d with $s < 0$ for certain values of the parameters (n, r, d) satisfying $n - d + r < 0$ follows immediately from the result for linear series with $s = 0$. We make this precise.

Corollary 4. *Let C be an arbitrary curve of genus g and let (n, r, d) be a triple such that $n - d + r < 0$, $n < g$, and $s < 0$. Then a general linear series $\mathcal{L}$ of degree d and dimension r does not admit a de Jonquières divisor of length n.*

Proof. Fix $r_1 := r - s$, such that $g - d + r_1 = g - d + r - s = 0$. Then we can conclude using Proposition 3 that the linear series does not admit de Jonquières divisors of length n if $n - d + r_1 < 0$, which is equivalent to the condition $n < g$. □

For non-special linear series we also immediately obtain that the space $DJ_n(\mathcal{L})$ is of expected non-negative dimension. This result will be useful at a later stage of the proof of Theorem 1 for special linear series, and is also interesting in its own right.

Proposition 5. *Let C be an arbitrary curve of genus g and let (n, r, d) be a triple such that $n - d + r \geq 0$ and $s = 0$. Then the space $DJ_n(\mathcal{L})$ is of expected dimension $n - d + r$. Moreover, if the dimension is zero, then a general linear series $\mathcal{L}$ of degree d and dimension r admits a finite number of de Jonquières divisor of length n.*

Proof. Consider the same Abel-Jacobi-type map φ as in the proof of Proposition 3. Here we have that $n \geq g$, hence the general fibre will indeed be of dimension $n - g = n - d + r$. If $n - d + r = 0$, from the finiteness of φ we conclude that the fibre of a general line bundle $L \in \mathrm{Pic}^d(C) = W^r_d(C)$ is finite. □

3.3. **The case of linear series of any dimension with $s = 1$.** The proof of the non-existence result in this case follows via residuation from the study of strata of abelian differentials in [Pol].

Proposition 6. *Let C be a general curve of genus g and let (n, r, d) be a triple such that $n - d + r < 0$ and $s = 1$. Then a general linear series $\mathcal{L}$ of degree d and dimension r does not admit a de Jonquières divisor of length n.*

Proof. Since $g - d + r = 1$, the residual linear series $K_C \otimes \mathcal{L}^\vee$ of $\mathcal{L}$ is of type g^0_{2g-2-d} i.e. it is given by an isolated divisor E of degree $2g - 2 - d$.

Since $|K_C|$ is complete, for any divisors D in $\mathcal{L}$, we can write $K_C \simeq \mathcal{O}(D+E)$.

Consider the subspace

$$\mathcal{P}_d := \{L \in \mathrm{Pic}^d(C) \mid h^1(L) = 1\} \subset \mathrm{Pic}^d(C).$$

It is easy to see that it has dimension $2g - 2 - d$.

Now let $\mu = (a_1, \ldots, a_n)$ with $a_i > 0$ be a partition of $2g - 2$. Consider the space

$$\mathcal{Q}(\mu) := \{(E; p_1, \ldots, p_n) \in C_{2g-d-2} \times C_n \mid \mathcal{O}(E + a_1 p_1 + \ldots + a_n p_n) \simeq K_C\}.$$

Polishchuk [Pol] shows that this space has dimension $n + g - d - 1$. Now take the following Abel-Jacobi type map

$$\varphi : \mathcal{Q}(\mu) \to \mathcal{P}_d$$
$$(E; p_1, \ldots, p_n) \mapsto \mathcal{O}(K_C - E).$$

Then $\dim \mathrm{im}\, \varphi \leq \dim \mathcal{Q}(\mu) = n + g - d - 1$. Moreover, note that $\dim \mathrm{im}\, \varphi < \dim \mathcal{P}$ if and only if $n < g - 1$. However this coincides exactly with our hypothesis $n < d - r = g - 1$. Thus, as in the proof of the previous lemma, the general linear series $\mathcal{L}$ with $s = 1$ does not admit a de Jonquières divisor of length n. □

3.4. **Degeneration argument for linear series of dimension $r \geq 2$.** In order to use degeneration techniques, we must first define our moduli space. Denote by $\mathcal{H}_{d,g,r}$ the unique component of the Hilbert scheme $\mathrm{Hilb}_{d,g,r}$ of curves of degree d and genus g in $\mathbb{P}^r$ which dominates the moduli space $\mathcal{M}_g$ of curves of genus g. Note that we shall refer to a curve lying in the component $\mathcal{H}_{d,g,r}$ as a curve with general moduli. Let $\mathcal{C} \to \mathcal{H}_{d,g,r}$ be the universal family and $\mathcal{C}^n$ be the n-th fibre product of $\mathcal{C}$ over $\mathcal{H}_{d,g,r}$.

As in Section 2, we define $\Sigma \subset \mathcal{C}^n$ as the degeneracy locus corresponding to collections $(C, p_1, \ldots, p_n)$, where the points $p_1, \ldots, p_n \in C$ are such that

$p_1 + \ldots + p_n \in DJ_n(\mathcal{L})$, and where $\mathcal{L}$ is the linear series embedding C in $\mathbb{P}^r$. In other words, it is the locus in C^n where

$$H^0(X, \mathcal{O}_X(1)(-a_1p_1 - \ldots - a_np_n)) \neq 0$$

for some positive integers $a_1, \ldots, a_n$ with $a_1 + \ldots + a_n = d$. Note that this also gives a reasonable definition for de Jonquières divisors on singular curves. We have the commutative diagram

$$\begin{array}{ccc} \Sigma & \xrightarrow{\iota} & C^n \\ & \searrow^{\pi} & \downarrow \\ & & \mathcal{H}_{d,g,r} \end{array}$$

where ι is the closed immersion of Σ into C^n and π is the (proper) map restricting the fibre product map $C^n \to \mathcal{H}_{d,g,r}$ to Σ.

The idea is to find nodal curves in $\mathcal{H}_{d,g,r}$ whose corresponding fibre in Σ is empty when $n - d + r < 0$. By the upper semicontinuity of the fibre dimension of π, we then conclude that a smooth curve with general moduli also has an empty fibre, i.e. its embedding does not admit de Jonquières divisors of length n when $n - d + r < 0$.

3.4.1. *The case of linear series of dimension $r = 2$.* Here the general $\mathcal{L}$ corresponds to a nodal plane model of the general curve C.

Proposition 7. *Let C be a general curve of genus g and let $(n, 2, d)$ be a triple such that $n - d + 2 < 0$ and $s > 1$ or $s < 0$. Then a general linear series $\mathcal{L}$ of type g^2_d does not admit a de Jonquières divisor of length n.*

Proof. The proof has three parts. The first part deals with the case $n - d + 2 = -1$, by rephrasing the argument in terms of multitangency conditions of a line in the plane with respect to C. Note that the argument here is independent of the genus, so both situations $s > 1$ and $s < 0$ are treated together. The rest of the proof deals with the case $n - d + 2 < -1$: in the second part we proceed by induction on the genus g of C and the degree d and dimension r of $\mathcal{L}$, with base case the linear series with $s = 1$. The purpose of the induction is to increase s at each step. The third part is again an argument by induction whereby s decreases at each step. The advantage of the proof by induction is the fact that all possible partitions of length n of d corresponding to the different de Jonquières divisors can be treated at once, rather than by a case by case analysis of different tangency conditions.

§1. Proof for $n - d + 2 = -1$.

If $d = 3$, we are in the case of the elliptic normal curve discussed in the introduction. If $d = 4$, then $n = 1$, implying that the de Jonquières divisor is of the form $4p$, which contradicts the result of [EH] mentioned at the beginning of the section.

For $d \geq 5$, the condition $n = d - 3$ translates into the curve admitting tangents through hyperflexes, bitangents through a flex point, or tritangents,

or in other words partitions of the type

$$4p_1 + p_2 + p_3 + \ldots + p_n,$$
$$3p_1 + 2p_2 + p_3 + \ldots + p_n,$$
$$2p_1 + 2p_2 + 2p_3 + p_4 + \ldots + p_n.$$

The first type of partition implies non-ordinary ramification of the linear series at p_1 and is immediately excluded, as discussed in the introduction.

We now discuss the case of the second type of partition. Let $\mathbb{P}^N = \mathbb{P}\,H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d))$ be the space parametrising all plane curves of degree d, where

$$N = \binom{d+2}{2} - 1.$$

Denote by $\mathcal{Z}^k$ the $(k+2)$-dimensional closed subscheme of the k-th symmetric power of $\mathbb{P}^2$ corresponding to collinear length k zero-cycles in $\mathbb{P}^2$ and let $\mathcal{Z}^0$ be the space of length one zero-cycles. Define the closed immersion $i : \mathcal{Z}^n \hookrightarrow \mathcal{Z}^d$ by

$$p_1 + p_2 + \ldots + p_n \mapsto 3p_1 + 2p_2 + p_3 + \ldots + p_n.$$

Let the morphism $\phi : \mathbb{P}^N \times (\mathbb{P}^2)^\vee \to \mathcal{Z}^d$ be given by the intersection product in $\mathcal{Z}^d$. Since the points $p_1, \ldots, p_n$ (with multiplicitiy) impose d independent conditions on $\mathbb{P}^N$, we have that

$$\dim \phi^{-1}(i(\mathcal{Z}^n)) = (N - d) + (n + 2) = N - 1.$$

Hence the canonical projection of $\phi^{-1}(i(\mathcal{Z}^n))$ to $\mathbb{P}^N$ is a closed subscheme $\mathbb{P}^N$ of dimension at most $N - 1$. We conclude that its complementary open set consists of the general plane curves of degree d that do not admit these partitions.

To deal with the remaining partitions, define the $(n + 2)$-dimensional space

$$\mathcal{Z} := \{(p_1 + p_2 + p_3, p_4, \ldots, p_n) \mid p_4, \ldots, p_n \in \mathcal{Z}^0 \text{ collinear to } p_1 + p_2 + p_3 \in \mathcal{Z}^3\}.$$

We again have a closed immersion $i : \mathcal{Z} \hookrightarrow \mathcal{Z}^d$ defined by

$$(p_1 + p_2 + p_3, p_4, \ldots, p_n) \mapsto 2p_1 + 2p_2 + 2p_3 + p_4 + \ldots + p_n.$$

Define as before the morphism $\phi : \mathbb{P}^N \times (\mathbb{P}^2)^\vee \to \mathcal{Z}^d$. As before, the canonical projection of $\phi^{-1}(i(\mathcal{Z}))$ to $\mathbb{P}^N$ is a closed subscheme $\mathbb{P}^N$ of dimension at most $N - 1$ and we are done.

§2. Proof for $s > 1$ and $n - d + 2 < -1$.

We start with a curve C of genus $g' = g - 2$ with a plane nodal model $\Gamma \subset \mathbb{P}^2$ of degree $d' = d - 1$ given by a g^2_{d-1} with $s = g' - d' + 2 \geq 1$ for which the statement of the proposition is true. In other words, we start with a g^2_{d-1} that does not admit de Jonquières divisors of length n' such that

$$n' - (d - 1) + 2 < 0.$$

Following [Ser], let γ be a general line on $\mathbb{P}^2$, so that

$$\Gamma \cap \gamma = \{q_1, \ldots, q_d\}.$$

Then we take C' to be the stable curve obtained by desingularising all multiple points of $\Gamma \cup \gamma$ except q_1, q_2, q_3. The curve C' now has arithmetic genus

$$p_a(C') = g' + 2 = g,$$

degree

$$\deg(C') = d' + 1 = d,$$

and furthermore

$$p_a(C') - \deg(C') + 2 = g - d + 2 = (g' - d' + 2) + 1 = s + 1.$$

Therefore by attaching γ to C we have increased s by one, which is what we were after.

Moreover, we have in fact a complete g^2_d on C'. By construction

$$h^0(C', \mathcal{O}_{C'}(1)) \geq 3.$$

Twisting the exact sequence

$$0 \to \mathcal{O}_\gamma(-q_1 - q_2 - q_3) \to \mathcal{O}_{C'} \to \mathcal{O}_C \to 0 \tag{1}$$

yields the long exact sequence of cohomology

$$0 \to H^0\left(\mathcal{O}_\gamma(1)(-q_1 - q_2 - q_3)\right) \to H^0\left(\mathcal{O}_{C'}(1)\right) \to H^0\left(\mathcal{O}_C(1)\right) \to \dots$$

from which we deduce that

$$h^0\left(\mathcal{O}_{C'}(1)\right) \leq h^0\left(\mathcal{O}_C(1)\right) = 3.$$

Therefore $h^0\left(\mathcal{O}_{C'}(1)\right) = 3$ and the g^2_d is complete.

Assume now towards a contradiction that a de Jonquières divisor of length n exists for this g^2_d, i.e. that

$$H^0(C', \mathcal{O}_{C'}(1)(-a_1p_1 - \dots - a_np_n)) \neq 0.$$

Let $D := a_1p_1 + \dots + a_np_n$, $D_\gamma := D \cdot \gamma$ and $D_C := D \cdot C$. We have the Mayer-Vietoris sequence

$$0 \to H^0(C', \mathcal{O}_{C'}(1)(-D)) \to H^0(C, \mathcal{O}_C(1)(-D_C)) \oplus H^0(\gamma, \mathcal{O}_\gamma(1)(-D_\gamma)) \to \mathbb{C}^3,$$

where the last map is given by

$$(u, v) \mapsto (u(q_1) - v(q_1), u(q_2) - v(q_2), u(q_3) - v(q_3)).$$

We have a few possibilities:

i) If both u and v are non-trivial sections then D is the pullback of the divisor given by the intersection of a hyperplane H in $\mathbb{P}^2$ with $\Gamma \cup \gamma$. In other words $D_\gamma = H \cdot \gamma = p_n$, say, and $D_C = H \cdot \Gamma = D - p_n$. If H does not pass through any of the nodes of $\Gamma \cup \gamma$, then the g^2_{d-1} corresponding to C admits a de Jonquières divisor of length $n - 1$. However

$$(n - 1) - (d - 1) + 2 = n - d + 2 < -1,$$

so we arrive at a contradiction with the induction hypothesis on C. See also Figure 1. On the other hand, if H passes through one of the nodes of $\Gamma \cup \gamma$, then the g^2_{d-1} on C admits a de Jonquières divisor of length n. However

$$n - (d - 1) + 2 = (n - d + 2) - 1 < 0,$$

which again contradicts the induction hypothesis on C. We note here that this case is the reason why we need a separate argument in the case $n - d + 2 = -1$. More precisely, if this is the case, then

$$n - (d-1) + 2 = (n - d + 2) + 1 = 0,$$

when we actually expect to find a finite number of de Jonquières divisors of length n, and hence we do not obtain a contradiction.

ii) If $u \equiv 0$, then C would be a line in $\mathbb{P}^2$, so we can discard this case.

iii) If $v \equiv 0$, then $H = \gamma$, and $u(q_1) = u(q_2) = u(q_3) = 0$ i.e. $u \in H^0(C, \mathcal{O}_C(1)(-D_C - q_1 - q_2 - q_3))$. Since u cannot be trivial, $\deg(D_C + q_1 + q_2 + q_3) \leq d - 1$. However, since $\deg D_C = \deg(\gamma \cdot \Gamma) = d - 1$, we must have that the q_i-s are in the support of D_C. Therefore $\deg(D_\gamma) = 1$, hence one of the q_i-s must be in its support. We continue in the same way as in i), in particular as in the case when H passes through a node.



FIGURE 1. Induction step for $r = 2$.

As a final step, we need to ensure that the induction step yields a curve with general moduli. Following [Ser], let $\mathcal{V}_{d,g}$ be the Severi variety parametrising plane irreducible nodal curves of degree d and genus g. The starting curve Γ is parametrised by a general point of an irreducible component of $\mathcal{V}_{d',g'}$ with the expected number of moduli. More precisely, the following two conditions are satisfied

i) lines in $\mathbb{P}^2$ cut the complete linear system $\mathcal{L} = |L|$ on the normalisation C of Γ

ii) the Petri map

$$\mu_0 : H^0(L) \otimes H^0(\omega_C \otimes L^\vee) \to H^0(\omega_C)$$

given by the multiplication of sections is injective.

These are sufficient conditions for the existence of irreducible components of $\mathcal{V}_{d,g}$ with general moduli. The aim now is to check that the inductive step preserves these properties. To begin with, note that $\mathcal{O}_{C'}(1)$ is determined by the pullback of a divisor cut by a line section on $\Gamma \cup \gamma$. One can check

(cf. [Ser] Proposition 2.3) that $\mu_0(\mathcal{O}_{C'}(1))$ is also injective. From Severi's classical theory we have a family of plane nodal curves of degree d

$$\begin{array}{ccc} \mathcal{X} & \subset & \mathbb{P}^2 \times S \\ \pi \downarrow & & \\ S & & \end{array}$$

parametrised by a smooth curve S, with central fibre

$$\mathcal{X}_0 = \Gamma \cup \gamma \subset \mathbb{P}^2$$

corresponding to the point $0 \in S$. Moreover all fibres $\mathcal{X}_s$ corresponding to $s \in S$ and $s \neq 0$ are irreducible nodal curves of genus g, so that

$$[\mathcal{X}_s] \in \mathcal{V}_{d,g}$$

and all their nodes specialise to the nodes of $\Gamma \cup \gamma$, with the exception of q_1, q_2, q_3.

Consider the desingularisation diagram

$$\begin{array}{ccccc} \mathcal{X}' & \xrightarrow{\nu} & \mathcal{X} & \hookrightarrow & \mathbb{P}^2 \times S \\ & \searrow & \downarrow \pi & & \\ & & S & & \end{array}$$

where ν is the blow-up of $\mathcal{X}$ along the closure of the singular locus of $\sigma^{-1}(S \setminus \{0\})$. By construction the fibre over $0 \in S$ is

$$\nu_0 : C' \to \Gamma \cup \gamma \subset \mathbb{P}^2 \times S$$

and for $s \neq 0$,

$$\nu_s : \mathcal{X}'_s \to \mathcal{X}_s \subset \mathbb{P}^2$$

is the normalisation map. By semicontinuity, we deduce that there is a C'' parametrised by a point in $\mathcal{V}_{d,g}$ satisfying the two properties above, since a linear system pulled back via ν_s over a sufficiently general $s \in S$ will have the same properties as the linear system $|\mathcal{O}_{C'}(1)|$ over C'. Hence, again by semicontinuity, there is an irreducible component with expected dimension of $\mathcal{V}_{d,g}$ and satisfying the two conditions above.

Finally we check that any general curve can be reached via this procedure, i.e we must check that

$$\rho(g, 2, d) \geq 0 \Rightarrow \rho(g', 2, d') \geq 0.$$

This is indeed the case:

$$\begin{aligned} \rho(g', 2, d') &= \rho(g-2, 2, d-1) \\ &= \rho(g, 2, d) + 1 > 0. \end{aligned} \tag{2}$$

§3. Proof for $s < 0$ and $n - d + 2 < -1$.

The argument is analogous to the one for $s > 1$. We start with a curve C of genus g with a plane nodal model $\Gamma \subset \mathbb{P}^2$ of degree $d' = d - 1$ given by a g^2_{d-1} with $s = g - d' + 2 \leq 0$ for which the statement of the proposition is

true. In other words, we start with a g^2_{d-1} that does not admit de Jonquières divisors of length n such that

$$n - (d-1) + 2 < 0.$$

Following [Ser], let γ be a general line on $\mathbb{P}^2$, so that

$$\Gamma \cap \gamma = \{q_1, \ldots, q_d\}.$$

Then we take C' to be the curve obtained by desingularising all multiple points of $\Gamma \cup \gamma$ except q_1. The curve C' now has arithmetic genus $p_a(C') = g$, degree $\deg(C') = d' + 1 = d$, and moreover

$$p_a(C') - \deg(C') + 2 = g - d + 2 = (g - d' + 2) - 1.$$

Thus by attaching γ to C we have decreased s by one, which is what we were after. The rest of the proof then proceeds just like before. □

3.4.2. *The case of linear series of dimension $r = 3$.* For $r \geq 3$, since the linear series $\mathcal{L}$ is general, it has no base points and its associated map

$$C \xrightarrow{|V|} \mathbb{P}^r$$

is an embedding (cf. [EH]). We denote the image of the embedding also by C. The proof of the non-existence result for $r = 3$ is similar to the one for $r = 2$, the main difference being the need for a smoothing argument.

Proposition 8. *Let C be a general curve of genus g and let $(n, 3, d)$ be a triple such that $n - d + 3 < 0$ and $s > 1$ or $s < 0$. Then a general linear series $\mathcal{L}$ of type g^3_d does not admit a de Jonquières divisor of length n.*

Proof. §1. Proof for $s > 1$.

The inductive step is the same as in the proof of Proposition 7. We start with a curve $C \subset \mathbb{P}^3$ of degree $d' = d - 1$, genus $g' = g - 2$, and with $s = g' - d' + 3 \geq 1$ which satisfies the proposition. Since there is a one-dimensional family of trisecant lines on C, we pick a line $\gamma \subset \mathbb{P}^3$ such that the intersection

$$C \cap \gamma = \{q_1, q_2, q_3\}$$

is transverse. As before, $X := C \cup \gamma$ has arithmetic genus $p_a(X) = g' + 2 = g$, degree $\deg(X) = d' + 1 = d$, and moreover $g - d + 3 = s + 1$, as desired. Finally, the exact sequence (1) yields analogously that $h^0(\mathcal{O}_X(1)) = 4$, i.e. the embedding is complete.

We now proceed as in the proof of Proposition 7, using the Mayer-Vietoris sequence

$$0 \to H^0(X, \mathcal{O}_X(1)(-D)) \to H^0(C, \mathcal{O}_C(1)(-D_C)) \oplus H^0(\gamma, \mathcal{O}_\gamma(1)(-D_\gamma)) \to \mathbb{C}_q,$$

where the last map is given by $(u, v) \mapsto u(q) - v(q)$, and moreover $D_\gamma = D \cdot \gamma$, and $D_C = D \cdot C$. The same case-by-case analysis as in the proof of Proposition 7 proves that $\mathcal{O}_X(1)$ does not admit a de Jonquières divisor of length n when $n - d + 3 < -1$.

However, we need to discuss the case $n-d+3=-1$ separately. Note that the possible partitions here are

$$\begin{aligned}
&5p_1+p_2+p_3+\ldots+p_n,\\
&4p_1+2p_2+p_3+\ldots+p_n,\\
&3p_1+3p_2+p_3+\ldots+p_n,\\
&3p_1+2p_2+2p_3+p_4+\ldots+p_n,\\
&2p_1+2p_2+2p_3+2p_4+p_5+\ldots p_n.
\end{aligned}$$

The first partition can be immediately excluded on grounds of ordinary ramification for general linear series on general curves. As mentioned before, if the hyperplane H passes through a node of X (i.e. the cases $u\equiv 0$ or $u(q)=v(q)=0$), then on C we have

$$n-(d-1)+3=0,$$

so we cannot use the induction hypothesis in the same way. However we can further degenerate X to a curve

$$X':=C'\cup\gamma_1\cup\ldots\cup\gamma_s,$$

where C' is a non-degenerate, non-special curve in $\mathbb{P}^3$ of degree $d-s$ and $\gamma_1,\ldots,\gamma_s$ are trisecant lines to C'. The Mayer-Vietoris exact sequence reads

$$\begin{aligned}
0\to H^0(X',\mathcal{O}_{X'}(1)(-D))\to\\
\to H^0(C',\mathcal{O}_{C'}(1)(-D_{C'}))\oplus H^0(\gamma_1,\mathcal{O}_{\gamma_1}(1)(-D_{\gamma_1}))\oplus\ldots\\
\ldots\oplus H^0(\gamma_s,\mathcal{O}_{\gamma_s}(1)(-D_{\gamma_s}))\to\mathbb{C}^{3s},
\end{aligned}$$

where $D_{C'}=D\cdot C'$, $D_{\gamma_i}=D\cdot\gamma_i$, and the last morphism is the usual one. A similar case-by-case analysis as in the proof of Proposition 7 and a comparison with the possible partitions that can occur reveal that the worst possibility is when the maximum number of sections $v_i\in H^0(\gamma_i,\mathcal{O}_{\gamma_i}(1)(-D_{\gamma_i}))$ vanish at nodes of X' belonging to different trisecant lines. For $s\leq 3$, this number is actually s, while for $s\geq 4$ it is equal to 4. This implies that the difference between the length and the degree of $D_{C'}$ equals s for $s\leq 3$ and it equals 4 for $s\geq 4$. It follows that C' is embedded by a g^3_{d-s} admitting a de Jonquières divisor of length n if $s\leq 4$, or $n-s+4$ otherwise. By Proposition 5, this g^3_{d-s} admits a $(s-1)$-dimensional family of de Jonquières divisors of length n, and a 3-dimensional family of de Jonquières divisors of length $n-s+4$. However, we have a one-dimensional family of trisecant lines to C', and for each degeneration we are free to choose s such trisecant lines. Hence we can choose them in such a way that they avoid the families of de Jonquières divisors and we are done.

To conclude the induction step we must smooth X to a curve with general moduli and a corresponding embedding with index of speciality $s+1$. We use the techniques of Sernesi [Ser] which we summarise below.

For a nodal curve $X\subset\mathbb{P}^r$ with $p_a(X)=g$ and $\deg(X)=d$, denote by T^1_X the sheaf defined through the exact sequence

$$0\to T_X\to T_{\mathbb{P}^r}|_X\to N_{X/\mathbb{P}^r}\to T^1_X\to 0.$$

Moreover, set

$$N'_X := \ker\left\{N_{X/\mathbb{P}^r} \to T^1_X\right\}.$$

Then, cf. [Ser], a sufficient condition for X to be flatly smoothable and for $\mathrm{Hilb}_{d,g,r}$ to be smooth and of expected dimension at the point $[X]$ is the vanishing

$$H^1(X, N'_X) = 0.$$

In our case, of course $r = 3$, $X = C \cup \gamma$, and C and γ meet transversely at three points q_1, q_2, q_3. Then we can write the following exact sequence on X

$$0 \to N_{\gamma/\mathbb{P}^3}(-q_1 - q_2 - q_3) \to N'_X \to N_{C/\mathbb{P}^3} \to 0.$$

This exact sequence implies that $H^1(N'_X) = 0$ if

$$H^1\left(C, N_{C/\mathbb{P}^3}\right) = H^1\left(\gamma, N_{\gamma/\mathbb{P}^3}(-q_1 - q_2 - q_3)\right) = 0.$$

But these conditions are satisfied: first of all, since we work with a Brill-Noether general curve C we have that indeed

$$H^1\left(C, N_{C/\mathbb{P}^3}\right) = 0.$$

Moreover, $N_{\gamma/\mathbb{P}^3} \simeq \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2}$, which means that

$$H^1\left(\gamma, N_{\gamma/\mathbb{P}^3}(-q_1 - q_2 - q_3)\right) = H^1\left(\mathcal{O}_{\mathbb{P}^1}(-2)\right)^{\oplus 2} = 0.$$

Arguing like in Proposition 2.3 of [Ser], we see that the Petri map corresponding to $\mathcal{O}_X(1)$ is injective and this guarantees that we can smooth X to a curve of general moduli. The smoothing of X' follows automatically.

The next step is to check that the induction step indeed increases the index of speciality for the general smooth curve. Denote by Z the irreducible component of expected dimension of $\mathrm{Hilb}_{d,g,3}$ containing X. Fix a general $\mathcal{X}_t \in Z$, which is smooth, as X is smoothable. Since X spans $\mathbb{P}^3$, $\mathcal{X}_t$ also spans $\mathbb{P}^3$ so that

$$h^0(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(1)) \geq 4.$$

Since $h^1(X, \mathcal{O}_X(1)) = s+1$, semicontinuity gives $h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(1)) \leq s+1$. On the other hand, Riemann-Roch yields

$$\begin{aligned} h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(1)) &= h^0(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(1)) - d + g - 1 \\ &\geq 3 - d + g = s + 1. \end{aligned}$$

Therefore

$$h^1(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(1)) = s + 1.$$

Finally, the same computation as in (2) shows that any Brill-Noether general curve may be reached via this induction step.

§2. Proof for $s < 0$.

Just like in the proof for $r = 2$, we start with a curve $C \subset \mathbb{P}^3$ of degree $d' = d - 1$, genus g, and $s = g - d' + 3 \leq 0$ for which the statement of the proposition is true. We attach a line γ at a point q, such that the intersection $C \cap \gamma$ is transverse. The new nodal curve $X := C \cup \gamma$ has arithmetic genus $p_a(X) = g$, degree $\deg(X) = d' + 1 = d$, and from Riemann-Roch $g - d + 3 = g - d' - 1 + 3 = s - 1$, as desired.

The same argument as in the proof of Proposition 7 shows that the embedding of X does not admit a de Jonquières divisor of length n. Note that in the problematic case $n - d + 3 = -1$, we simply degenerate X to

$$X' := C' \cup \gamma_1 \cup \ldots \cup \gamma_s,$$

where C' is a non-special curve of degree $d-s$ and the γ_i are lines intersecting C' transversely in one point. We then argue as in the proof for $s > 1$.

To conclude the induction step we must ensure that X can be smoothed in $\mathbb{P}^3$ to a curve with general moduli. As in the proof of Proposition 8 the vanishing of $H^1(X, N'_X)$ is given by the vanishing

$$H^1\left(C, N_{C/\,\mathbb{P}^3}\right) = H^1\left(\gamma, N_{\gamma/\,\mathbb{P}^3}(-q)\right) = 0.$$

But this follows immediately, and we conclude that X is flatly smoothable. Moreover, the Petri map is automatically injective for $s < 0$, hence X belongs to the unique irreducible component $\mathcal{H}_{d,g,3}$ of the Hilbert scheme, and this concludes the proof. □

3.4.3. *The case of linear series of dimension $r \geq 4$.* The proof for linear series with $s > 1$ or $s < 0$ for $r \geq 4$ proceeds by induction. Here we are no longer able to guarantee that the general curve C has a trisecant line, hence we cannot proceed by induction on s for $s > 1$ by attaching such a line to a starting curve in $\mathbb{P}^4$. Instead we construct an inductive step that increases r and d, keeps s constant, and uses as base case embedded space curves satisfying Proposition 8. For non-complete linear series, however, the induction step decreases s at each step by attaching a rational normal plane curve to C in an appropriate fashion. Luckily, the geometry of the inductive step in both cases is well-behaved in the situation of hyperplanes passing through nodes, therefore we need not treat the case $n - d + r = -1$ separately any longer.

Proposition 9. *Let C be a general curve of genus g and let (n, r, d) be a triple such that $n - d + r < 0$, $s > 1$ or $s < 0$, and $r \geq 4$. Then a general linear series $\mathcal{L}$ of type g^r_d does not admit a de Jonquières divisor of length n.*

Proof. §1. Proof for $s > 1$.

We begin with a curve $C \subset \mathbb{P}^{r-1} \subset \mathbb{P}^r$ of degree $d' = d-1$, genus g, and for which the statement of the proposition is true. Attach to C a line γ such that the intersection $\gamma \cap C = \{q\}$ is transverse and $\gamma \subset \mathbb{P}^r$, but is not contained in the same hyperplane as C. The resulting curve $X := C \cup \gamma$ has arithmetic genus $p_a(X) = g$, degree $\deg(X) = d' + 1 = d$, as shown in Figure 2.

An exact sequence analogous to (1) applied at each step shows that X is embedded by a complete linear series, i.e.

$$h^0(\mathcal{O}_X(1)) = r + 1,$$

and furthermore we have that $g - d + (r + 1) = g - d' + r = s$, i.e. the index of speciality of the embedding remains unchanged.

Assume towards a contradiction that the embedding of X admits a de Jonquières divisor of length n, i.e. that

$$H^0(X, \mathcal{O}_X(1)(-D)) \neq 0.$$



FIGURE 2. Construction for $r \geq 4$ and $s > 1$.

We have the Mayer-Vietoris sequence

$$0 \to H^0(X, \mathcal{O}_X(1)(-D)) \to H^0(C, \mathcal{O}_C(1)(-D_C)) \oplus H^0(\gamma, \mathcal{O}_\gamma(1)(-D_\gamma)) \to \mathbb{C}_q,$$

where the last map is given by $(u, v) \mapsto u(q) - v(q)$. As usual, let $D_C = D \cdot C$ and $D_\gamma = D \cdot \gamma$. We have a few cases

i) If u and v are non-trivial sections, D is given by the intersection of a hyperplane $H \in \mathbb{P}^r$ with X. If H does not pass through any of the nodes, then C is embedded by a g^{r-1}_{d-1} admitting a de Jonquières divisor of length $n-1$, which contradicts the induction hypothesis. If H does pass through the node, then the g^{r-1}_{d-1} admits a de Jonquières divisor of length n, which again contradicts the hypothesis.

ii) If $u \equiv 0$, then $C \subset H$ and $v(q) = 0$. Therefore $v \in H^0(\gamma, \mathcal{O}_\gamma(1)(-D_\gamma - q))$. Since v is non-trivial and $\deg(\mathcal{O}_\gamma(1)) = 1$, q must be one of the points in the support of D, in particular $D_\gamma = q$. Then $D_C = D - q$ and the induction step follows as in i), in particular as in the case where H passed through the node.

iii) If $v \equiv 0$, then $\gamma \subset H$ and $u(q) = 0$. Therefore $u \in H^0(C, \mathcal{O}_C(1)(-D_C - q))$. Since u is non-trivial, $\deg(D_C + q) \leq d - 1$. However, since $\deg D_C = \deg(H \cdot C) = d - 1$, q must be one of the points in the support of D_C. Then $\deg D_\gamma = 1$ and in particular $D_\gamma = q$. We proceed with the induction step as in the case of the hyperplane passing through the node from i).

We now prove that X smooths to a curve with general moduli. To do this we use certain nice components of $\mathrm{Hilb}_{\mathrm{d,g,r}}$ introduced in [BE]. To be precise, for fixed integers r, d, g such that $r \geq 3$, $g \geq 0$, there exists a unique irreducible component $W(d, g; r) \subset \mathrm{Hilb}_{\mathrm{d,g,r}}$ which is generically smooth and of expected dimension

$$3g - 3 + \rho(g, r, d)$$

such that a general $X \in W(d, g; r)$ has the following properties:

i) X is a smooth connected non-degenerate curve of degree d, genus g and such that $h^1(X, N_X) = 0$;
ii) if $d \geq g + r$, then $h^1(X, \mathcal{O}_X(1)) = 0$;
iii) if
$$d - r < g \leq d - r + \lfloor (d - r - 2)/(r - 2) \rfloor,$$
then X is linearly normal and $h^1(X, \mathcal{O}_X(2)) = 0$;
iv) if $\rho(g, r, d) \geq 0$, then C has general moduli, i.e.
$$W(d, g; r) = \mathcal{H}_{d,g,r};$$
v) if $\rho(g, r, d) < 0$, then the general fibre of the natural rational map
$$W(d, g; r) \dashrightarrow \mathcal{M}_g$$
has dimension given by $\dim(\mathrm{Aut}(\mathbb{P}^r))) = r^2 + 2r$, i.e. $W(d, g; r)$ has the right number moduli.

So we must prove that X is smoothable and that $X \in W(d, g; r)$. Our argument is based on tools developed in [BF] and [Ser].

First degenerate C in $\mathbb{P}^{r-1}$ to a general union
$$\Gamma := \Gamma_1 \cup C_1 \cup \ldots \cup C_s,$$
defined as follows: Γ_1 is a smooth connected curve of degree
$$d_1 := d' - (r - 1)s$$
and genus
$$g_1 := g - rs,$$
which imply $g_1 - d_1 + r - 1 = 0$, i.e.
$$h^1(\Gamma_1, \mathcal{O}_{\Gamma_1}(1)) = 0.$$
Next, the C_i are pairwise disjoint rational normal curves in $\mathbb{P}^{r-1}$, each intersecting Γ_1 in $r + 1$ points. Moving γ in a family of projectively equivalent lines, we may assume that γ intersects Γ_1 in one point, while $\gamma \cap C_i = \emptyset$, for all i. Moreover, an argument using the exact sequence (1) yields
$$H^1(\Gamma_1 \cup \gamma, \mathcal{O}_{\Gamma_1 \cup \gamma}) = 0.$$
Since Γ_1 is non-special and γ is a line, the Mayer-Vietoris exact sequence
$$0 \to N_{\Gamma_1 \cup \gamma} \to N_{\Gamma_1 \cup \gamma}|_{\Gamma_1} \oplus N_{\Gamma_1 \cup \gamma}|_{\gamma} \to N_{\Gamma_1 \cup \gamma}|_{\Gamma \cap \gamma} \to 0$$
yields
$$H^1(\Gamma_1 \cup \gamma, N_{\Gamma_1 \cup \gamma}) = 0,$$
i.e. $\Gamma_1 \cup \gamma$ is smoothable in $\mathbb{P}^r$. Moreover the fact that the embedding is non-special guarantees that $\Gamma_1 \cup \gamma \in \mathcal{H}_{d_1+1,g_1,r} = W(d_1 + 1, g_1; r)$. To finish the argument, we still need to attach the rational normal curves C_i. The generality of the resulting curve is ensured by a repeated application of Lemma 7 in [BF].

Hence we have showed that X is smoothable to a curve with general moduli in $\mathbb{P}^r$. The same argument by semicontinuity as in the proof of Proposition 8 shows that the index of speciality stays the same for the smooth curve with general moduli.

Finally, we again check that all general curves can be reached by this construction, i.e. that

$$\rho(g,r,d) \geq 0 \Rightarrow \rho(g,r-1,d-1) \geq 0.$$

As before, this follows immediately:

$$\begin{aligned}\rho(g,r-1,d-1) &= g - r(g-d+1+r-1)\\ &= g - r(g-d+r)\\ &= \rho(g,r,d) + s > 0.\end{aligned}$$

§2. Proof for $s < 0$.

In the case of non-complete linear series we begin instead with a curve $C \subset \mathbb{P}^{r-1} \subset \mathbb{P}^r$ of degree $d' = d-2$, genus g, and such that the statement of the proposition is true. Attach to C a curve γ of degree two such that the intersection $C \cap \gamma = q$ is transverse and such that γ is not contained in the same hyperplane as C. See also Figure 3. The curve $X := C \cup \gamma$ has arithmetic genus $p_a(X) = g$ and degree $\deg(X) = d' + 2 = d$. The same arguments as before show that X is embedded by a linear series with $g - d + r + 1 = g - d' + r - 1 = s - 1$, which is what we were after.



FIGURE 3. Construction for $r \geq 4$ and $s < 0$.

Assume again towards a contradiction that the embedding of X admits a de Jonquières divisor of length n. The degenerate cases follow as in the $s > 1$ case. We discuss only the non-degenerate ones, meaning that there exists some hyperplane H in $\mathbb{P}^r$ which does not contain any of the components of X, and which intersects X in d points with the appropriate multiplicities. There are three possibilities

i) H does not pass through q and it is tangent to γ at one point, say p_1. Then we are left with a de Jonquières divisor of degree $d-2$ and length $n-1$ on C. On the other hand,

$$(n-1) - (d-2) + (r-1) = n - d + r < 0,$$

so we arrive at a contradiction with the inductive hypothesis.

ii) H does not pass through q and intersects γ transversely at two points, say p_1 and p_2. Then we obtain a de Jonquières divisor of degree $d-2$ and length $n-2$ on C. However,

$$(n-2)-(d-2)+(r-1)=n-d+r-1<0,$$

and we again arrive at a contradiction.

iii) H passes through q and is tangent to γ at that point. We then follow the same line of reasoning as in point i).

iv) H passes through q and intersects γ in another distinct point, say p_1. Then we are left with a de Jonquières divisor of degree $d-2$ and length $n-1$. However,

$$(n-1)-(d-2)+(r-1)=n-d+r<0,$$

and we conclude by contradiction that X cannot be embedded by a g^r_d admitting a de Jonquières divisor of length n.

The smoothing to a curve with general moduli argument is analogous to the one shown in the proof for $s<1$. □

References

[ACG] E. Arbarello, M. Cornalba, and P.A. Griffiths. *Geometry of Algebraic Curves II*, volume 268 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2011.

[ACGH] E. Arbarello, M. Cornalba, P.A. Griffiths, and J. Harris. *Geometry of Algebraic Curves I*, volume 267 of *Grundlehren der mathematischen Wissenschaften*. Springer, 1985.

[BCG+] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Moeller. Compactification of strata of abelian differentials. arXiv:1604.08834 [math.AG].

[BE] E. Ballico and Ph. Ellia. On the existence of curves with maximal rank in $\mathbb{P}^n$. *Journal für die reine und angewandte Mathematik*, 397:1–22, 1989.

[BF] E. Ballico and C. Fontanari. Normally generated line bundles on general curves, II. *Journal of Pure and Applied Algebra*, 214(8):1450 – 1455, 2010.

[dJ] E. de Jonquières. Mémoire sur les contacts multiples d'ordre quelconque des courbes, etc. *Crelle*, 66:289–321, 1866.

[EH] D. Eisenbud and J. Harris. Divisors on general curves and cuspidal rational curves. *Inventiones mathematicae*, 74:371–418, 1983.

[FP] G. Farkas and R. Pandharipande. The moduli space of twisted canonical divisors. arXiv:1508.07940 [math.AG].

[Mac] I.G. MacDonald. Symmetric products of an algebraic curve. *Topology*, 1:319–343, 1962.

[Mat] A. Mattuck. Secant bundles on symmetric products. *American Journal of Mathematics*, 87:779–797, 1965.

[Pol] A. Polishchuk. Moduli spaces of curves with effective r-spin structures. In *Gromov-Witten theory of spin curves and orbifolds*, volume 403 of *Contemporary Mathematics*, pages 1–20. American Mathematical Society, 2006.

[Ser] E. Sernesi. On the existence of certain families of curves. *Inventiones mathematicae*, 75(1):25–57, 1984.

[Vai] I. Vainsencher. Counting divisors with prescribed singularities. *Transactions of the American Mathematical Society*, 267(2):399–422, 1981.

Humboldt-Universität zu Berlin, Institut für Mathematik, Unter den Linden 6, 10099 Berlin, Germany

E-mail address: ungurean@math.hu-berlin.de